



The New York Times |

SCIENCE

# Pacific Sea Otters' Failure to Thrive Confounds Wildlife Sleuths

By **INGFEI CHEN** FEB. 27, 2012

MONTEREY, Calif. — On a fog-shrouded morning in Monterey Bay, wildlife researchers are out to capture a southern sea otter named Blanca — part of a three-year project to learn why her species, hunted to near extinction a century ago, is still in trouble here despite decades of efforts to bring it back.

Blanca is not cooperating.

Because wild sea otters bolt at the whiff of human presence, the only way to catch one is when it is asleep. Blanca is tagged with a radio transmitter, and scientists onshore are tracking her by telemetry and telescope.

About 8:30 a.m., she begins diving for crabs in a kelp bed off Cannery Row. In a skiff on the bay, three otter biologists — Tim Tinker, Brian Hatfield and Joe Tomoleoni — wait for her to stop feeding and take a nap.

And wait. And wait. When Blanca finally dozes, five hours after the tracking began, Mr. Hatfield and Mr. Tomoleoni slip into the water 270 yards away, with scuba gear and underwater scooters rigged with nets. Long minutes pass.

Then Dr. Tinker, watching with binoculars from the boat, sees her awaken and

plunge beneath the surface. “She just saw you,” he tells his companions by radio. “Target is gone. It’s over.”

For the wildlife biologists, a clear explanation for the sea otters’ failure to thrive is proving just as elusive. Almost wiped out by fur traders, the species rebounded after an international ban on commercial otter hunting in 1911. But today, the otter population in California is just 2,700, in a mosaic of small, separate colonies off the coast, down from perhaps as many as 16,000 in the past.

Multiple factors are stalling the recovery. One popular view, supported by veterinary pathologists who study dead otters, primarily blames coastal pollution — in the form of parasites, bacteria, toxins and chemicals.

But Dr. Tinker and other biologists say that, at least in the areas where the sea otter population is highest, off Monterey and nearby Big Sur, the underlying problem is simply that the otters are running out of food.

While they are not starving to death, they are depleting their favorite prey, sea urchins and abalone, and having to spend more time hunting. Poor nutrition is compromising their fitness to survive diseases or other threats, said Dr. Tinker, who runs the United States Geological Survey’s otter research program. “They’re not getting enough food to make it through.” Reports from Dr. Tinker’s team also suggest that otters are particularly vulnerable to sharks.

Bridging the two scientific camps, Dr. Tinker is working closely with veterinary experts and biologists at the California Department of Fish and Game, the Monterey Bay Aquarium, the University of California and elsewhere. The wildlife sleuths have been tracking diet, behavior, diseases, births and deaths among 90 radio-tagged sea otters that live off urban Monterey or pristine Big Sur. The sites differ mainly in that Monterey Bay receives more polluted runoff.

Last fall, the team was recapturing the otters to take more blood samples, pluck whiskers and retrieve a small, implanted, pen-shaped instrument (all with anesthesia) from each. The instrument recorded a trove of data on body temperature and the time and depth of every dive an otter had made in the past year.

By telescope, the scientists observed what the otters ate in more than 20,000 foraging dives. More information is coming from a novel test analyzing the chemical composition of the otter whiskers (based on the principle “you are what you eat,” said Seth D. Newsome, a research collaborator from the University of Wyoming).

And a new genetic technique detects whether pollutants and pathogens are impairing the otters’ immunological health, even before they get sick. The new blood test screens activity in 14 key genes, said Lizabeth Bowen, a geneticist at the Geological Survey who developed the test with Jeffrey Stott of the University of California, Davis.

The genetic signatures can reveal whether an animal is experiencing subtle physiological stress, inflammation or infection by bacteria or parasites, Dr. Bowen said — or reacting to exposure to pollutants like PCBs. The testing cannot as yet tell precisely which contaminant may be stressing the otters.

The Big Sur-Monterey study is part of a larger, multiagency effort called the Pacific Nearshore Project, which is comparing nine distinct sea otter populations and the health of their coastal habitats in the northern Pacific.

The broader project is investigating why some colonies in southeastern Alaska, British Columbia and Washington that were growing rapidly two decades ago — by 20 percent a year — have seen that rate slowed by half, said its leader, James L. Bodkin of the Geological Survey’s Alaska Science Center. The California otters’ growth rate is even more lackluster: usually less than 5 percent a year and, lately, near zero.

Sea otters are remarkably voracious: To survive frigid waters, they must fuel a high metabolism by consuming 25 to 30 percent of their body weight every day. Veterinary scientists, who tend to favor the coastal-pollution explanation, note that the otters dine on many types of shellfish and invertebrates that are prone to accumulating contaminants.

As a result, the animals “are getting hit with so many things,” said Melissa A. Miller, a veterinary pathologist at the California Fish and Game Department who autopsies stranded otters. “I picture it sometimes almost like otters are sitting there

right at that land-sea interface with their mouths open.”

In 2010, Dr. Miller and her associates reported evidence that microcystin, a toxin from blue-green algae that live only in freshwater lakes and streams, had killed at least 21 sea otters. Another toxin, domoic acid, is also deadly to the animals. Such poisons are generated by harmful growths of algae that can be fed by fertilizers in agricultural runoff.

Dr. David Jessup, a veterinarian retired from the state wildlife agency, says other leading killers include disease-causing parasites transmitted in feces from cats and opossums; infections by bacteria in human or animal feces; and industrial pollutants, which may subtly affect otter immune defenses. These factors all “have some connection to human activities,” he said.

But to ecologists, emerging evidence instead strongly suggests that elevated rates of infectious diseases are mostly a symptom of a larger problem — insufficient food resources and malnutrition.

“When animals reach a point of extreme nutritional stress,” Dr. Tinker said, “they will succumb to whatever particular stressor they encounter first” — whether a parasite or toxin, a boat strike or a shark attack that a well-fed otter might otherwise fend off or evade.

Even without harmful pathogens from land, “I think they would be dying from something else,” said James Estes, a marine ecologist at the University of California, Santa Cruz.

While these ecologists do not minimize the importance of cleaning up coastal pollution, they doubt it would lead to a major rebound in the sea otter population off Big Sur and Monterey.

Dr. Tinker says these areas probably cannot sustain any more otters, given the available supplies of sea urchins, abalone and other shellfish.

“Their rate of food acquisition has declined to a point where they’re pretty much spending as much time feeding as they can,” he said — 40 to 50 percent of each day. They are scrawny compared with the “big, round happy otters” at San Nicolas Island

in Southern California, where their prey abounds and the otters forage only 25 percent of the time.

The results from the current comparison study, due this spring, may help resolve some of the debate by answering whether otters fare better in the cleaner waters at Big Sur.

Preliminary genetic tests indicate higher stress, inflammation and exposure to pathogens and pollutants in the Monterey group, Dr. Tinker said. But survival data have not been analyzed.

For now, the research partners agree to disagree. No single answer can explain the California sea otter mystery. For instance, food is abundant elsewhere in the otter range, so why aren't animals rapidly multiplying there and moving into unoccupied territories?

No one knows. The reason may be a high rate of shark attacks in some places, or a land-based pathogen in others, Dr. Tinker said. And, because those low-density otter colonies still have much growth potential, he said, combating pollution might still be the best long-term answer to the sea otters' plight.

A version of this article appears in print on February 28, 2012, on page D3 of the New York edition with the headline: Pacific Sea Otters' Failure to Thrive Confounds Wildlife Sleuths.